

FOOD ALLERGY ACTION PLAN

Name: _____ Date of birth: _____

Allergy to: _____

Weight: _____ lbs. Asthma: _____ Yes (higher risk for a severe reaction) _____ No

Extremely reactive to the following foods: _____

THEREFORE:

_____, if checked, give epinephrine for ANY symptoms if the allergen was *likely* eaten.

_____, if checked give epinephrine immediately if the allergen was *definitely* eaten, even if no symptoms noted.

Any severe symptoms after suspected or known ingestion:

One or more of the following:

Lung: Short of breath, wheeze, repetitive cough

Heart: Pale, blue, faint, weak pulse, dizzy, confused

Throat: Tight, hoarse, trouble breathing/swallowing

Mouth: Obstructive swelling (tongue and/or lips)

Skin: Many hives over body

Or *combination of symptoms* from different body areas:

Skin: Hives, itchy rashes, swelling (e.g., eyes, lips)

Gut: Vomiting, crampy pain

PLAN

1. **INJECT EPINEPHRINE IMMEDIATELY**
2. Call 911
3. Begin monitoring
4. Give additional medications: *
 - Antihistamine
 - Inhaler (bronchodilator) if asthmatic

*Antihistamines & inhalers/brochodilators are not to be depended upon to treat a severe reaction (anaphylaxis). **USE EPINEPHRINE**

Mild symptoms only:

Mouth: Itchy mouth

Skin: A few hives around mouth/face, mild itch

GUT: Mild nausea/discomfort

PLAN

1. **GIVE ANTIHISTAMINE**
2. Stay with student: alert healthcare professionals and parent
3. IF symptoms progress (see above), **USE EPINEPHRINE**
4. Begin monitoring

Medications/Doses

Epinephrine (brand and dose): _____

Antihistamine (brand and dose): _____

Other (e.g., inhaler-bronchodilator if asthmatic): _____

Monitoring

Stay with the student; alert healthcare professionals and the parent. Tell rescue squad epinephrine was given; request an ambulance with epinephrine. Note time when epinephrine was administered. A second dose of epinephrine can be given 5 minutes or more after the first if symptoms persist or recur. For a severe reaction, consider keeping student lying on back with legs raised. Treat student even if parents cannot be reached.

Parent /Guardian Signature Date

Physician/Healthcare Provider Signature Date

A food allergy response kit should contain at least two doses of epinephrine, other medications as noted by the student's physician, and a copy of this Food Allergy Action Plan.

A kit must accompany the student if he/she is off school grounds (i.e., field trip).

Contacts

Call 911

Physician: _____ Phone: _____

Parent/Guardian: _____ Phone: _____

Other emergency contacts

Name/relationship: _____ Phone: _____

Name /relationship: _____ Phone: _____

Reference: FAAN (www.foodallergy.org)